

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**  
Cambridge Ordinary Level

**MARK SCHEME for the October/November 2014 series**

**4024 MATHEMATICS (SYLLABUS D)**

**4024/11**

Paper 1, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

<b>Page 2</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge O Level – October/November 2014</b>	<b>4024</b>	<b>11</b>

### Abbreviations

cao	correct answer only
cso	correct solution only
dep	dependent
ft	follow through after error
isw	ignore subsequent working
oe	or equivalent
SC	Special Case
www	without wrong working
soi	seen or implied

Question	Answers	Mark	Part marks
<b>1 (a)</b>	41 006	1	
<b>(b)</b>	240 000	1	
<b>2 (a)</b>	12	1	
<b>(b)</b>	(0).08	1	
<b>3 (a)</b>	$\frac{3}{100}$ cao	1	
<b>(b)</b>	82	1	
<b>4 (a)</b>	64	1	
<b>(b)</b>	67	1	
<b>5</b>	$(2a - 3b)(c + 2d)$	2	<b>B1</b> for one of the partial factorisations $c(2a - 3b)$ ; $2d(2a - 3b)$ ; $2a(c + 2d)$ ; $-3b(c + 2d)$ or their negatives, seen.
<b>6 (a)</b>	$\frac{8}{9}$	1	
<b>(b)</b>	28	1	
<b>(c)</b>	90	1	
<b>7</b>	A correct method to eliminate one variable Either $x = 4$ or $y = -1$ WWW. Both $x = 4$ and $y = -1$ WWW.	M1 A1 A1	If [0] earned, then award <b>C1</b> for a pair of values that satisfy either equation.

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2014	4024	11

8	(a)	9	1	
	(b)	8	1	
	(c)	25	1	
9		8 WWW	3	<b>M1</b> for a recognisable attempt at Pythagoras' Theorem with sides 10 and 6. <b>M1</b> for $(AT^2 =) 10^2 - 6^2$ oe
10	(a)	$P \cap Q \cap R'$ oe	1	
	(b)	47	2	<b>M1</b> for Cricket set inside the Football set, e.g. in a Venn diagram; Ans. = $30+8+9$ ; "30 play both cricket and football".
11	(a)	$\begin{pmatrix} 330 \\ 417 \end{pmatrix}$	2	<b>B1</b> for 330 or 417 in a (2 by 1) matrix, or for (330 417).
	(b)	P shows the amount earned in Week 1 and Week 2, oe	1 dep	Must refer to (i) the amount earned (money, earnings, \$, etc) and (ii) the two weeks.
12	(a)	930	1	
	(b)	$\frac{2s - an}{n}$ oe	2	<b>M1</b> for correct first step, e.g. $2s = an + bn$ ; $s = na/2 + nb/2$ or <b>B1</b> for a correct expression for $b$ seen in working, but followed by an error.
13		$d = \frac{5v^2}{64}$		<b>M1</b> for $d = kv^2$ , or for $5 = k \times 64$ ;
		125	3	<b>B1</b> for $k = 5/64$ , or for $\frac{d}{5} = \frac{40^2}{8^2}$
14	(a)	3.65	1	
	(b)	60 WWW	3	<b>B1</b> for 192; or for cost price = \$120, so i by (profit =) \$72. <b>M1</b> for $\left(\frac{\text{their}192 - \text{their}120}{\text{their}120}\right) \times 100$ oe
15	(a)	Triangle $ABC$ drawn with an acceptable $C$ .	2	<b>B1</b> for $AC = 7$ cm or <b>B1</b> for $\angle CAB = 130^\circ$
	(b)	21 to 22 inclusive, WWW; Or FT their triangle, provided the perp. height is not one of the sides, WWW.	2✓	<b>M1</b> for $\frac{1}{2}$ base $\times$ height with matching base and height.

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2014	4024	11

16	(a)	$x + y = 6$ drawn correctly	1	<b>B1</b> for <b>R</b> correctly bordered by the lines $y = 2$ and $x = -1$ ; or FT appropriate shading between their sloping lines, provided one is correct	
	(b)	$2y + x = 4$ drawn correctly	1		
	(c)	Correct region shaded, (FT for sloping lines with one correct line).	2✓		
17	(a)	Valid method, with $\frac{1}{2}(11+7) \times 4 \times 5$ oe, leading to 180	1	<b>AG</b>	
	(b)	20 WWW	3	<b>B1</b> for 22 500 or 0.18 and <b>M1</b> for $\sqrt[3]{\frac{\text{figs } 225}{\text{figs } 18}}$ soi	
18	(a)	14 41	1		
	(b)	149	1		
	(c)	(i)	2 5 10 17		1
		(ii)	$n^2 - 1$ oe		1
19	(a)	$1.36 \times 10^9$	1	<b>B1</b> for figs 793, or for $N \times 10^5$ with $1 < N < 10$ .	
	(b)	(i)	$5.6 \times 10^9$		1
		(ii)	$7.93 \times 10^5$		2
20	(a)	F	1		
	(b)	C	1		
	(c)	B	1		
	(d)	E	1		
21	(a)	(i)	... alternate (angles) ...	1	<b>M1</b> for $\frac{180 - 58}{2}$ , or <b>B1</b> for a base angle = $61^\circ$
		(ii)	$119^\circ$	2	
	(b)	120 WWW	2	<b>C1</b> for 240. <b>M1</b> for $2x + 80 + 95 + 125 = 540$ , oe	

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2014	4024	11

22	(a)	42	1	
	(b)	Correct plots at 20, 40, 60, 90, 120 and CF curve drawn	2	<b>B1</b> for three or four correct plots
	(c)	(i) 62 to 64 inclusive (ii) 41 to 46 inclusive WWW, FT ( $F_{80}-F_{50}$ ) from their graph.	1✓ 2✓	FT from their CF graph <b>M1</b> for attempt to calculate ( $F_{80}-F_{50}$ ) from their graph.
23	(a)	(i) the point <i>B</i> marked correctly	1	If [0] scored in (a), in (aiii) award <b>B1</b> for the vector $\begin{pmatrix} -6 \\ 1 \end{pmatrix}$ soi.
		(ii) the point <i>C</i> marked correctly	1	
		(iii) the point <i>D</i> marked correctly	1	
	(b)	(i) $\mathbf{q} - \mathbf{p}$	1	<b>M1</b> for $\mathbf{OT} = \mathbf{OR} + \mathbf{RT}$ Or for $\mathbf{OT} = \mathbf{OP} + \mathbf{PR} + \mathbf{RT}$ Or for $\mathbf{OT} = \mathbf{OQ} + \mathbf{QR} + \mathbf{RT}$ Or equivalents in terms of $\mathbf{p}$ and $\mathbf{q}$ .
		(ii) $\frac{2}{3}\mathbf{p} + \frac{1}{3}\mathbf{q}$	1✓	
		(iii) $\frac{1}{3}\mathbf{q} - \left(\frac{4}{3}\right)\mathbf{p}$ , or FT their(ii) $-2\mathbf{p}$	2✓	